

Telemedicine: Can In-Person Pre-treatment Communication be Expanded by Video Consultation?

S. Guhl¹ · L. Linngrön¹ · B. Rosenberg¹  · N. Hosten¹ · M. Kirsch¹

Received: 19 August 2019 / Accepted: 30 August 2019
© The Author(s) 2019

To the Editor,

Informed consent for radiological or other interventions should give a patient sufficient time to make an informed decision. Currently, patients typically have to be present, in-person, to be briefed about procedures and an extra appointment is often necessary. While in an urban setting this is mostly just a nuisance, in a rural area, similar to ours, it may not be possible at all for patients with limited access to transport. In some countries, teleconsultation via videoconference has proved beneficial in comparable situations [1].

We performed a pilot study, approved by the ethical committee of our university. Fifty patients were 1:1 randomized and one group was briefed face-to-face, the other via videoconferencing. With our hospital being located in a rural area, patients travel 50.2 km to our department for periradicular therapy (mean: range 1–110 km). Thirty-two percent of study patients already used videoconferencing Apps such as Skype[®] or Facetime[®] in their private lives. Patients provided written consent firstly to the intervention itself and secondly to the pilot study presented here. They were informed that participation in the study was voluntary. Patients were referred for CT-guided periradicular or facet joint infiltration aimed at reducing chronic back pain. Groups did not show significant differences regarding the age ($t(48) = -1.827, p = 0.074, n = 50$) or

distribution between the sexes ($\chi^2(1) = 0.89, p = 0.765, n = 50$) (57 years vs. 64 years and 68% males vs. 61% males for videoconference vs. in-person, respectively). Interventions were performed by a radiologist supported by technical assistants. Both groups received the same pre-treatment discussion by the performing radiologist [2], followed by the legally required face-to-face briefing for the videoconferencing group later. To compare the effectiveness of both options, a questionnaire was read to patients in a telephone call by a blinded study nurse 24 h after the briefing [3]. A total of eight questions covered atmosphere and necessity of information, quality of the doctor/patient relationship and the feeling of being taken seriously by the physician. The remaining questions covered the items summarized in Figs. 1 and 2.

Patients who received the pre-treatment briefing by videoconference remembered significantly more (Mann-Whitney U test: $U = 210.000, p = 0.038, r = 0.2932$) of the mentioned side effects compared to patients who received the pre-treatment briefing in-person (Fig. 1). Further, the recall of radiation exposure was significantly higher when communicated in a videoconference ($\chi^2(1) = 3.947, p = 0.047, n = 50, \phi = -0.281$, Fig. 2). For patient satisfaction with pre-treatment communication and the other variables related to knowledge acquisition, no significant differences emerged.

Preoperative discussion by videoconferencing was equal to, or better than, face-to-face discussion. We assume that patients easily focus on a monitor, and distraction is thus reduced [4]. While there may be extra costs to cover the equipment, the process of informing patients about procedure may actually be facilitated (less logistical effort, patients may be given a specific time window for the call). Briefings via videoconference could be saved, with

S. Guhl and L. Linngrön contributed equally to this study.

✉ B. Rosenberg
britta.rosenberg1@uni-greifswald.de

¹ Department of Diagnostic Radiology and Neuroradiology, Universitätsmedizin Greifswald, Ferdinand-Sauerbruch-Straße, 17475 Greifswald, Germany




Fig. 1 Recall was higher for videoconference briefings than for face-to-face briefings




Fig. 2 Recall of items explained in briefings

additional viewings made available to patients. Additional information material may also be provided easily.

In our opinion, it is worthwhile to evaluate “informed consent to treatment in low population density areas by teleconsultation” in larger studies. A higher number of participants are necessary as effect sizes were small; more realistic scenarios with clinic to home videoconferencing should be employed; improving the process of blinding of study nurses, if possible, as patients tended to mention the videoconference in interviews; legal aspects (saving the interviews digitally; use of electronic devices for getting

informed consent in one study group) must be clarified beforehand, and endpoints of studies should be chosen in a way that validated questionnaires can be used.

Funding This study was funded by Grant INT-08-0001 of INTERREG IV A of the European Union.

Compliance with Ethical Standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical Approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards.

Informed Consent Informed consent was obtained from all individual participants included in the study.

Consent for Publication For this type of study, consent for publication is not required.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Bradford N, Caffery L, Smith A. Telehealth services in rural and remote Australia: a systematic review of models of care and factors influencing success and sustainability. *Rural Remote Health*. 2016;16:3808.
2. Maurer MH, Disch AC, Hartwig T, Renz DM, Schreiter N, Hamm B, Streitparth F. Outcome study of real-time MR-guided cervical periradicular injection therapy in an open 1.0 Tesla MRI system. *Cardiovasc Intervent Radiol*. 2014;37:756–62. <https://doi.org/10.1007/s00270-013-0738-4>.
3. O'Dwyer H, Lyon S, Fotheringham T, Lee M. Informed consent for interventional radiology procedures: a survey detailing current European practice. *Cardiovasc Intervent Radiol*. 2003;26:428–33. <https://doi.org/10.1007/s00270-003-0058-1>.
4. Tschamper MK, Jakobsen R. Parents' experiences of videoconference as a tool for multidisciplinary information exchange for children with epilepsy and disability. *J Clin Nurs*. 2019;28:1506–16. <https://doi.org/10.1111/jocn.14755>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.